arXiv:1210.3209v1 [hep-ph] 11 Oct 2012

Charmonium dissociation and recombination at LHC: Revisiting comovers

E.G. Ferreiro[*]

Departamento de Física de Partículas and IGFAE, Universidade de Santiago de Compostela,
15782 Santiago de Compostela, Spain

Abstract

Charmonium production at heavy-ion colliders is considered within the comover interaction model. The formalism includes both comover dissociation of J/ψ's and possible secondary J/ψ production through recombination. The estimation of this effect is made without involving free parameters. The comover interaction model also incorporates a comprehensive treatment of initial-state nuclear shadowing. With these tools, the model properly describes the centrality, transverse momentum and rapidity dependence of experimental data from PbPb collisions at the LHC energy of $\sqrt{s} = 2.76$ TeV. We present predictions for PbPb collisions at $\sqrt{s} = 5.5$ TeV.

---

[*] *E-mail address:* elena@fpaxp1.usc.es

1 Introduction

Lattice QCD calculations predict that, at sufficiently large energy densities, hadronic matter undergoes a phase transition to a plasma of deconfined quarks and gluons (QGP). Substantial activity has been dedicated to the research of high-energy heavy-ion collisions in order to reveal the existence of this phase transition and to analyze the properties of strongly interacting matter in the new phase. The study of quarkonium production and suppression is among the most interesting investigations in this field since, in the presence of a QGP, the charmonium yield would be further suppressed due to color Debye screening [1]. Indeed, such an anomalous suppression was first observed in PbPb collisions at top CERN SPS energy [2]. Alternatively, the SPS experimental results could also be described in terms of final state interactions of the $c\bar{c}$ pairs with the dense medium created in the collision, the so-called comover interaction model (CIM) [3, 4, 5, 6]. This model does not assume thermal equilibrium and, thus, does not use thermodynamical concepts.

The theoretical extrapolations to RHIC and LHC energies were guided mainly by two trends. On the one hand, models assuming a deconfined phase during the collision pointed to the growing importance of secondary J/ψ production due to regeneration of $c\bar{c}$ pairs in the plasma. The total amount of $c\bar{c}$ pairs is assumed to be created in hard interactions during the early stages of the collision. Then, either using kinetic theory and solving rate equations for the subsequent dissociation and recombination of charmonium [7, 8, 9], or assuming statistical coalescence at freeze-out [10, 11, 12], one obtains the final J/ψ yield. On the other hand, the CIM with only dissociation of J/ψ's predicted [13] a stronger suppression at RHIC than at SPS due to a larger density of produced soft particles in the collision. It also predicted a stronger suppression at mid-rapidity –where the comover density is maximal– than at forward rapidities. Nevertheless, measurements of J/ψ production in AuAu collisions at RHIC, $\sqrt{s} = 200$ GeV, gave surprising results: the suppression at mid-rapidity was on the same level as at SPS [14, 15]. Furthermore, the suppression at forward rapidity in AuAu collisions was stronger than at mid-rapidity.

These facts showed that in the CIM, which is based on the well-known gain and loss differential equations in transport theory, the introduction of a recombination term is actually required for a detailed balance [16]. In the present work we use the updated version of the CIM [16], that allows recombination of $c\bar{c}$ pairs into secondary J/ψ's. We will estimate this effect using the density of charm in proton-proton collisions at the same energy. Therefore, the model does not involve any additional parameters. In Section 2 we present the details of the model; the effects related to the initial-state shadowing and nuclear absorption, together with comover dissociation and regeneration are described. In Section 3 we present our results for PbPb collisions at the LHC energy of $\sqrt{s} = 2.76$ TeV and predictions at $\sqrt{s} = 5.5$ TeV. Conclusions and final remarks are given in Section 4.

2 Description of the model

In this section, we give a short and updated description of the CIM. The present version of this model contains a comprehensive treatment of initial-state nuclear effects, such as nuclear shadowing, together with nuclear absorption, interaction with the co-moving matter and the recombination of $c\bar{c}$ into secondary J/ψ's.

Nuclear effects in nucleus-nucleus collisions are usually expressed through the so-called nuclear modification factor, $R_{AB}^{J/\psi}(b)$, defined as the ratio of the J/ψ yield in AA and pp scaled by the number of binary nucleon-nucleon collisions, $n(b)$. We have then

$$
\begin{aligned}
R_{AB}^{J/\psi}(b) &= \frac{\mathrm{d}N_{AB}^{J/\psi}/\mathrm{d}y}{n(b)\,\mathrm{d}N_{pp}^{J/\psi}/\mathrm{d}y} \\
&= \frac{\int \mathrm{d}^2 s\, \sigma_{AB}(b)\, n(b,s)\, S^{abs}(b,s)\, S^{sh}_{J/\psi}(b,s)\, S^{co}(b,s)}{\int \mathrm{d}^2 s\, \sigma_{AB}(b)\, n(b,s)}\ , \qquad (1)
\end{aligned}
$$

where $\sigma_{AB}(b) = 1 - \exp[-\sigma_{pp}\, AB\, T_{AB}(b)]$, $T_{AB}(b) = \int \mathrm{d}^2 s T_A(s) T_B(b-s)$ is the nuclear overlapping function and $T_A(b)$ is obtained from Woods-Saxon nuclear densities [17]. In eq. (1),

$$
n(b,s) \ = \ \sigma_{pp} AB\, T_A(s)\, T_B(b-s)/\sigma_{AB}(b)\ , \qquad (2)
$$

where upon integration over $\mathrm{d}^2 s$ we obtain the number of binary nucleon-nucleon collisions at impact parameter b, $n(b)$.

The three additional factors in the numerator of eq. (1), S^{abs}, S^{sh} and S^{co}, denote the effects of nuclear absorption, shadowing and interaction with the co-moving matter –both dissociation and recombination–, respectively. They will be defined below.

2.1 Nuclear absorption

At low energies the primordial spectrum of particles created in scattering off a nucleus is mainly altered by interactions with the nuclear matter –spectators nucleons– they traverse on the way out to the detector. This effect is called nuclear absorption and is usually parameterized within a probabilistic Glauber model. One has

$$
S^{abs}(b,s) = \frac{[1 - \exp(-A\ T_A(s)\ \sigma_{abs})][1 - \exp(-B\ T_B(b-s)\sigma_{abs})]}{\sigma^2_{abs}\ AB\ T_A(s)\ T_B(b-s)}\ . \qquad (3)
$$

Several novel features are expected to appear at high energies due to the change in the space-time picture of particle production and the coherence effects [18]. In this way, the break-up cross section becomes $\sigma_{abs} = 0$ at high energies [19, 20, 21]. From now on, we will take $S^{abs} = 1$ for LHC energies, which implies a negligible nuclear absorption.

2.2 Shadowing

Coherence effects will lead to nuclear shadowing for both soft and hard processes at high energy, and therefore also for the production of heavy flavor. Shadowing can be calculated within the Glauber-Gribov theory [22] making use of the generalized Schwimmer model of multiple scattering [23]. In this case the second suppression factor in eq. (1) is given by

$$S^{sh}(b,s,y) \;=\; \frac{1}{1\,+\,AF(y_A)T_A(s)}\,\frac{1}{1\,+\,BF(y_B)T_B(b-s)}\ , \tag{4}$$

where the function $F(y)$ encodes the dynamics of shadowing. Following the spirit of the model presented in [24, 25], where shadowing corrections are given without free parameters in terms of the triple-Pomeron coupling determined from diffraction data, one can write:

$$F(y) = 4\pi \int_{y_{min}}^{y_{max}} dy\ \frac{1}{\sigma_P(y)}\ \frac{d\sigma^{PPP}}{dydt}\bigg|_{t=0} \;=\; C\left[\exp\left(\Delta y_{max}\right)\,-\,\exp\left(\Delta y_{min}\right)\right]/\Delta\ , \tag{5}$$

It represents the coherence effects due to shadowing corrections expressed as the ratio of the triple-Pomeron cross section over the single-Pomeron exchange. We take $y_{min} = \ell n(R_A m_N/\sqrt{3})$, where m_N is the nucleon mass and $R_A = 0.82\ A^{1/3} + 0.58$ fm - the gaussian nuclear radius. The value of y_{max} depends on the rapidity of the produced particle, $y_{max} = \ln(s/m_T^2)/2 \pm y$ with the $+(-)$ sign if the particle is produced in the hemisphere of nucleus $B(A)$. m_T is the transverse mass of the produced particle. We have used $\Delta = 0.13$ and $C = 0.04$ fm^2 ($C/\Delta = 0.31$ fm^2) corresponding to the Pomeron intercept $\alpha_P(0) = 1.13$. With the shadowing resulting from the above equations a good description of the centrality dependence of charged multiplicities is obtained both at RHIC [26] and LHC [27] energies.

The energy dependence of these different effects can be summarized as follows. Particle production at SPS is dominated by low-energy effects, i.e. nuclear absorption and small nuclear shadowing, while RHIC already belongs to the high-energy regime, where nuclear shadowing becomes relevant, and the combined effect of shadowing and energy-momentum conservation should be accounted for at forward rapidities. At LHC, shadowing is expected to be strong while nuclear absorption is a small effect and can be neglected. We will now proceed with the discussion of comover interaction effects.

2.3 Dissociation by comover interaction and recombination

The CIM was developed to explain both the suppression of charmonium yields [3, 4, 5, 6, 13, 28, 29] and the strangeness enhancement [30, 31] in nucleus-nucleus collisions at the SPS. Neglecting possible recombination effects, the rate equation governing the density of charmonium in the final state, $N_{J/\psi}$, can be written in a simple form assuming a pure longitudinal expansion of the system and boost invariance. For an AA collision the density of J/ψ at a given transverse coordinate, s, impact parameter b, and rapidity y is given by

$$\tau \frac{\mathrm{d}N_{J/\psi}}{\mathrm{d}\tau}\,(b,s,y) \;=\; -\sigma_{co} N^{co}(b,s,y) N_{J/\psi}(b,s,y)\;, \tag{6}$$

where σ_{co} is the cross section of charmonium dissociation due to interactions with the co-moving medium, with density N^{co}. It is found from fits to low-energy experimental data to be $\sigma_{co} = 0.65$ mb [5]. To incorporate the effects of recombination, we have to include an additional gain term proportional to the squared density of open charm produced in the collision. Then eq. (6) is generalized to

$$\tau \frac{\mathrm{d}N_{J/\psi}}{\mathrm{d}\tau}\,(b,s,y) \;=\; -\sigma_{co}\left[N^{co}(b,s,y) N_{J/\psi}(b,s,y) \;-\; N_c(b,s,y) N_{\bar{c}}(b,s,y)\right]\;, \tag{7}$$

where we have assumed that the effective recombination cross section is equal to the dissociation cross section*. This extension of the model therefore does not involve additional parameters.

Equation (7) cannot be solved analytically. We approximate its solution as

$$S^{co}(b,s,y) \;=\; \exp\left\{-\sigma_{co}\left[N^{co}(b,s,y) \;-\; \frac{N_c(b,s,y) N_{\bar{c}}(b,s,y)}{N_{J/\psi}(b,s,y)}\right] \ln\left[\frac{N^{co}(b,s,y)}{N_{pp}(0)}\right]\right\}\;, \tag{8}$$

resembling the exact solution of eq. (6), since the first term in the exponent of eq. (8) is exactly the survival probability of a J/ψ interacting with comovers [13]. The density of comovers is calculated following the same lines as in [27] together with the proper shadowing correction. Then

$$N^{co}(b,s,y) \;=\; N^{co}_{NS}(b,s,y)\, S^{sh}_{ch}(b,s,y)\;, \tag{9}$$

where S^{sh}_{ch} denotes the shadowing for light particles. The non-shadowed (NS) multiplicity of comovers is taken as proportional to the number of nucleon-nucleon collision according to eq. (2)

$$N^{co}_{NS}(b,s,y) \;=\; N^{pp}(b,s,y)\, n(b,s)\;, \tag{10}$$

where N^{pp} represents the comover density in pp collisions, essentially $N^{pp}(y) = \frac{3}{2}(\mathrm{d}N^{ch}/\mathrm{d}y)^{pp}$. In fact, when using at mid-rapidity the value of $(\mathrm{d}N^{ch}/\mathrm{d}y)^{pp}_{y=0} = 3.9$, compatible with the inelastic value of $dN^{pp}/d\eta$ at $\sqrt{s} = 2.76$ TeV, a good agreement with experimental data on charged particle multiplicities is obtained [27].

The density of open and hidden charm in AA collisions, $N_c, N_{\bar{c}}$ and $N_{J/\psi}$, respectively, can be computed from their densities in pp collisions as $N^{AA}_c(b,s) = n(b,s) S^{sh}_{HQ}(b,s) N^{pp}_c$, with similar expression for $N^{AA}_{\bar{c}}$ and $N^{AA}_{J/\psi}$. Here $n(b,s)$ is given by eq. (2) and S^{sh}_{HQ} is the shadowing factor for heavy quark production, given by eq. (4). Then eq. (8) becomes

$$S^{co}(b,s,y) \;=\; \exp\left\{-\sigma_{co}\left[N^{co}(b,s,y) - C(y) n(b,s) S^{sh}_{HQ}(b,s)\right] \ln\left[\frac{N^{co}(b,s,y)}{N_{pp}(0)}\right]\right\} \tag{11}$$

---

[*] These two cross-sections have to be similar but not necessarily equal. We have taken the simplest possibility.

where

$$C(y) \;=\; \frac{\left(\mathrm{d}N_{pp}^{c\bar{c}}/\mathrm{d}y\right)^2}{\mathrm{d}N_{pp}^{J/\psi}/\mathrm{d}y} \;=\; \frac{\left(\mathrm{d}\sigma_{pp}^{c\bar{c}}/\mathrm{d}y\right)^2}{\sigma_{pp}\,\mathrm{d}\sigma_{pp}^{J/\psi}/\mathrm{d}y}\,. \tag{12}$$

The quantities in the rightmost term in eq. (12) are all related to pp collisions at the corresponding energy. The value for $\mathrm{d}\sigma_{pp}^{J/\psi}/\mathrm{d}y$ can be taken from ALICE [32] or a model for extrapolation of the experimental results [33]. The $c\bar{c}$ pairs are mostly in charmed mesons, such as D and D^* and the corresponding values could be extracted from the experiment [34], leading to an estimation of $\mathrm{d}\sigma_{pp}^{c\bar{c}}/\mathrm{d}y$, as we will discuss below. For σ_{pp} we use the non-diffractive value σ_{pp}=54 mb [27].

With σ_{co} fixed from experiments at low energy, where recombination effects are negligible, the model, formulated above, should be self-consistent at high energies[†].

3 Results

Our results for the centrality dependence of the J/ψ nuclear modification factor in PbPb collisions at 2.76 TeV are presented in Fig. 1 compared to ALICE experimental data [35, 36, 37, 38] at mid and forward rapidities. The different contributions to J/ψ suppression are shown. Note that the initial-state effect is just the shadowing, which can induce a suppression of $R_{AA} = 0.6$ for the more central collisions [39]. The combined effect of shadowing and comover dissociation gives a too strong suppression compared to experimental data. We therefore proceed to estimate the effect of recombination.

For the charmonium cross section pp measurements around mid-rapidity are available from ALICE [32] at 2.76 TeV which corresponds to $\frac{d\sigma_{pp}^{J/\psi}}{dy} = 3.73\ \mu$b. We consider that realistic values of C at mid-rapidity at 2.76 TeV are in the range of a minimum value of 2 up to a maximal value of 3 which corresponds to a cross section $\frac{d\sigma_{pp}^{c\bar{c}}}{dy} \approx 0.6 \div 0.8$ mb. This agrees with the estimated values in [40], and corresponds to a $\sigma_{c\bar{c}}^{tot}$ around 5 mb, which agrees well with experimental data [34]. These values are higher than the ones reported in [41], where data is also reproduced. Note nevertheless that there is no contradiction, since in [41] the initial-state shadowing is not considered. This shadowing, that affects also heavy flavours, would imply an extra suppression leading naturally to the choice of higher input charm cross sections $\mathrm{d}\sigma_{pp}^{c\bar{c}}/\mathrm{d}y$. In other words, one can consider that in the approach developed in [41] the choice of smaller $\frac{d\sigma_{pp}^{c\bar{c}}}{dy}$, $\frac{d\sigma_{pp}^{c\bar{c}}}{dy} \approx 0.3 \div 0.4$ mb, takes into account an effect of shadowing that reduces the input charm cross section up to 1/2.

We expect the effect of recombination to be stronger at mid than at forward rapidities. At $y \neq 0$ the recombination term is smaller since the rapidity distribution of D, D^* is narrower than the one of

---

[†] Note, however, that σ_{co} could change when the energy increases. We do not expect this effect to be important and, since we are unable to evaluate the magnitude of this eventual change, we shall use the same value $\sigma_{co} = 0.65$ mb at all energies.



Figure 1: Results on the centrality dependence of the J/ψ nuclear modification in PbPb collisions at 2.76 TeV at mid (left) and forward (right) rapidity compared to ALICE data [35, 36, 37, 38]. The dashed line corresponds to the **shadowing** effect. The lowest line corresponds to the combined effect of **shadowing + comover dissociation**. The shadowed area corresponds to our result when **shadowing + comover dissociation + recombination** are taken into account. The uncertainty takes into account the variation between the minimum and maximal value of C.

comovers, which induces a decrease of the C-value. This will produce a decrease of $R_{AA}^{J/\psi}$ with increasing y. Note nevertheless that this effect may be compensated by the increase of $R_{AA}^{J/\psi}$ due to a smaller density of comovers at $y \neq 0$, which induces less dissociation. We have chosen a smooth behaviour of $C(y)$ with rapidity, which in the rapidity range $2.5 < y < 4$ corresponds to a mean reduction in the C-value of the order of 20%. Taking $\frac{d\sigma_{pp}^{J/\psi}}{dy}$ in this forward rapidity range, $\frac{d\sigma_{pp}^{J/\psi}}{dy} = 2.23$ μb [32], these C-values correspond to an input charm cross section $\frac{d\sigma_{pp}^{c\bar{c}}}{dy} \approx 0.4 \div 0.6$ mb in the forward rapidity region. Our procedure gives a reasonable description of data when recombination is taken into account.

Some considerations proceed here:

First, the behaviour of the nuclear modification factor $R_{AA}^{J/\psi}$ for different rapidity ranges changes depending of the amount of recombination considered. If the recombination C-value is considered to be small –lower limit–, the total suppression will be controlled essentially by the comover dissociation, and the nuclear modification factor will follow the same behaviour as the usual comover suppression, but with a higher absolute value. On the other hand, when the C-value is taken as its upper limit, the recombination term controls the total suppression, inducing a decrease of the nuclear modification factor when going to forward rapidities.

Second, the shadowing corrections for heavy quarks we use here have been calculated in a very simple way, according to eq. (4) within the Glauber-Gribov theory. The advantage is that all calculations can be easily done analytically. This shadowing roughly agrees with EKS98/nDSg predictions [39, 42, 43, 44, 45], in particular in the mid-rapidity region. In Fig. 2 we show the comparison of the shadowing model applied here with some recent calculations developed within the framework of Refs. [44, 39]. One can see that the Glauber-Gribov inspired model agrees with the lowest band of nDSg according to [44, 39]. On the other side, calculations of the shadowing developed within the framework of the Color Evaporation model [45], lead in general to an slightly higher suppression than [44]. Because of this we consider that the shadowing presented here agrees with the models mentioned above within uncertainties. Nevertheless, the Glauber-Gribov shadowing is almost constant with rapidity. Even if the y-dependence of the different shadowing models in AA collisions and in the rapidity ranges here considered is quite flat, as it shown in [35] and Refs. [39, 45] therein, we are aware that our approach can induce an small overestimation of the shadowing suppression in the forward rapidity region. In fact we contemplate the upgrading of the CIM by the introduction of EKS98/nDSg according to [39, 44].



Figure 2: Centrality dependence of the Glauber-Gribov shadowing (eq. (4)) compared to EKS98/nDSg calculations performed according to [44, 39] in PbPb collisions at $\sqrt{s} = 2.76$ TeV.

We proceed now to study the behaviour of the nuclear modification factor when different transverse momentum cuts are introduced. In Fig. 3 (left) our results for 2 different p_T ranges, $0 < p_T < 2$ GeV and $5 < p_T < 8$ GeV, are compared to ALICE forward rapidity data [38]. Clearly, the amount of recombination is more important in the low p_T region. The comparison to CMS data at mid-rapidity and higher p_T, $6.5 < p_T < 30$ GeV [47, 48, 49], in Fig. 3 (right) emphasizes this finding: the amount of recombination is much smaller than in the mid-rapidity range at low p_T, as shown in Fig. 1 (left). It is important to remark that this experimental fact –stronger suppression at higher p_T– can not be due to initial-state shadowing effects: the presence of shadowing corrections, more relevant at lower p_T, conspirates in the opposite direction.



Figure 3: Results on the centrality dependence of the J/ψ nuclear modification factor in PbPb collisions at 2.76 TeV compared to ALICE forward rapidity data [38] in the p_T ranges $0 < p_T < 2$ GeV and $5 < p_T < 8$ GeV (left) and compared to CMS mid-rapidity data [47, 48, 49] in the p_T range $6.5 < p_T < 30$ GeV (right). The dashed line corresponds to the **shadowing** effect. The lowest line (right) corresponds to the combined effect of **shadowing + comover dissociation**. The shadowed area corresponds to our result when **shadowing + comover dissociation + recombination** are taken into account. The uncertainty takes into account the variation between the minimum and maximal value of C.

Moreover, our results on the dependence of the J/ψ nuclear modification factor on the transverse momentum are shown in Fig. 4 compared to ALICE [37] and CMS [47, 48] data. An overall agreement is obtained.



Figure 4: Results on the transverse dependence of the J/ψ nuclear modification in PbPb collisions at 2.76 TeV compared to CMS data [47, 48] at mid-rapidity (left) and to ALICE [38] and CMS data [47] at forward rapidity (left). The CMS data on the left part of the figure corresponds to the rapidity range $1.6 < |y| < 2.4$, while the ALICE data lie in the range $2.5 < y < 4$. The dashed line corresponds to the **shadowing** effect. The lowest line corresponds to the combined effect of **shadowing + comover dissociation**. The shadowed area corresponds to our result when **shadowing + comover dissociation + recombination** are taken into account. The uncertainty takes into account the variation between the minimum and maximal value of C.

We continue by showing our results versus rapidity. In Fig. 5 (left) we compare our results with CMS data [47, 48] in the mid-rapidity region. An overall agreement is obtained. The fact that we do not reproduce the detailed behaviour of ALICE data [37], as can be seen in Fig. 5 (right), is extremely instructive. As we have mentioned above, we have chosen a very conservative behaviour of $C(y)$ with rapidity, which leads to a mean reduction in the C-value of the order of 20% in the rapidity range $2.5 < y < 4$. The data in this region agrees with this choice in the interval $2.5 < y < 3.5$, while for the most forward points an scenario with smaller amount of recombination or only dissociation without recombination is clearly favoured.



Figure 5: Results on the rapidity dependence of the J/ψ nuclear modification in PbPb collisions at 2.76 TeV compared to CMS data [47, 48] at mid-rapidity (left) and to ALICE [37] at mid and forward rapidity (left). The lowest line corresponds to the combined effect of **shadowing + comover dissociation**. The shadowed area corresponds to our result when **shadowing + comover dissociation + recombination** are taken into account. The uncertainty takes into account the variation between the minimum and maximal value of C.

We finish by presenting our predictions for the LHC energy of 5.5 TeV. Here we have let vary our C-value between 2 and 4. Taking $BR_{ll} \times \frac{d\sigma_{pp}^{J/\psi}}{dy}|_{y=0} \approx 350$ nb [33] and $\sigma_{pp} = 60$ mb, these C-values correspond to an input charm cross section $\frac{d\sigma_{pp}^{c\bar{c}}}{dy} \approx 0.8 \div 1.15$ mb in the mid-rapidity region. In the rapidity range $2.5 < y < 4$ we take a mean reduction in the C-value of the order of 20%, which corresponds to an input charm cross section $\frac{d\sigma_{pp}^{c\bar{c}}}{dy} \approx 0.6 \approx 0.9$ mb when $BR_{ll} \times \frac{d\sigma_{pp}^{J/\psi}}{dy} \approx 250$ nb [33]. Our results in the mid and forward rapidity ranges are plotted in Fig. 6

Note that in the previous work [16] stronger shadowing corrections –of around 20% larger than the present ones– for the heavy quark production were considered, which implied both less recombination effects and stronger total suppression. This demonstrates that the implementation of the initial-state effects is more relevant than what is currently assumed in the recombination approaches, since it affects the probability of regeneration.



Figure 6: Results on the centrality dependence of the J/ψ nuclear modification in PbPb collisions at 5.5 TeV at mid (left) and forward (right) rapidity. The dashed line corresponds to the **shadowing** effect. The lowest line corresponds to the combined effect of **shadowing + comover dissociation**. The shadowed area corresponds to our result when **shadowing + comover dissociation + recombination** are taken into account. The uncertainty takes into account the variation between the minimum and maximal value of C. For the mid-rapidity region, we have taken, from down to up, $\frac{d\sigma_{pp}^{c\bar{c}}}{dy} = 0.8, 0.85, 0.9, 0.95, 1.0, 1.1, 1.15$ mb and for the forward rapidity region $\frac{d\sigma_{pp}^{c\bar{c}}}{dy} = 0.65, 0.67, 0.71, 0.74, 0.82, 0.84, 0.87, 0.89$ mb.

4 Conclusion

In this work we have studied the combined effect of J/ψ dissociation and recombination of $c\bar{c}$ pairs into J/ψ in the comover interaction model. This model does not assume thermal equilibrium of the matter produced in the collision and includes a comprehensive treatment of initial-state effects, such as shadowing. We estimate the magnitude of the recombination term from J/ψ and open charm yields in pp collisions at LHC. Without any adjustable parameters, the centrality, transverse momentum and rapidity dependence of experimental data is reproduced.

In our approach, the magnitude of the recombination effect is controlled by the total charm cross section in pp collisions. Note that, contrary to the results in [16], where the combined effect of initial-state shadowing and comover dissociation appeared to overcome the effect of parton recombination at 5.5 TeV, we find here that the recombination effects are of crucial importance in PbPb collisions at LHC energies and can dominate over the suppression, in agreement with [41, 40, 46]. The reason for this dissent is the fact that a different approach for the shadowing factor for heavy quark production was used in [16], which minimalized the amount of recombination and led to an overestimation of the total suppression.

Also, as a general remark, we are aware that the comover interaction model at these energies should not be considered to describe a final-state interaction at the hadronic level. Indeed, at small values of the proper time these comovers should be considered as a dense partonic medium. A large contribution to the comover interaction comes from the few first fm/c, where the system is in partonic or pre-hadronic stage.

Acknowledgments

It is a pleasure to thank R. Arnaldi, A. Capella, F. Fleuret, J.-P. Lansberg and E. Scomparin for useful exchanges. This work was partially supported by the Ministerio de Ciencia (Spain) & the IN2P3 (France) (AIC- D-2011-0740).

References

[1] T. Matsui, H. Satz, Phys. Lett. B **178**, 416 (1986)

[2] B. Alessandro, et al., Eur. Phys. J. C **39**, 335 (2005)

[3] A. Capella, A. Kaidalov, A. Kouider Akil, C. Gerschel, Phys. Lett. B **393**, 431 (1997)

[4] N. Armesto, A. Capella, Phys. Lett. B **430**, 23 (1998)

[5] N. Armesto, A. Capella, E.G. Ferreiro, Phys. Rev. C **59**, 395 (1999)

[6] A. Capella, E.G. Ferreiro, A.B. Kaidalov, Phys. Rev. Lett. **85**, 2080 (2000)

[7] R. Thews, M. Schroedter, J. Rafelski, Phys. Rev. C **63**, 054905 (2001)

[8] L. Grandchamp, R. Rapp, Nucl. Phys. A **709**, 415 (2002)

[9] L. Yan, P. Zhuang, N. Xu, Phys. Rev. Lett. **97**, 232301 (2006)

[10] P. Braun-Munzinger, J. Stachel, Phys. Lett. B **490**, 196 (2000)

[11] A. Andronic, P. Braun-Munzinger, K. Redlich, J. Stachel, Phys. Lett. B **571**, 36 (2003)

[12] A.P. Kostyuk, M.I. Gorenstein, H. Stöcker, W. Greiner, Phys. Rev. C **68**, 041902 (2003)

[13] A. Capella, E.G. Ferreiro, Eur. Phys. J. C **42**, 419 (2005)

[14] A. Adare, et al., Phys. Rev. Lett. **98**, 232301 (2007)

[15] M.J. Leitch, J. Phys. G **34**, S453 (2007)

[16] A. Capella, L. Bravina, E. G. Ferreiro, A. B. Kaidalov, K. Tywoniuk and E. Zabrodin, Eur. Phys. J. C **58**, 437 (2008)

[17] C.W. De Jager, H. De Vries and C. De Vries, Atom. Data Nucl. Data Tabl. **14**, 479 (1974)

[18] M.A. Braun, C. Pajares, C.A. Salgado, N. Armesto, A. Capella, Nucl. Phys. B **509**, 357 (1998)

[19] A. Capella and E. G. Ferreiro, Phys. Rev. C **76**, 064906 (2007)

[20] I. C. Arsene, L. Bravina, A. B. Kaidalov, K. Tywoniuk and E. Zabrodin, Phys. Lett. B **660**, 176 (2008)

[21] K. Tywoniuk, I. Arsene, L. V. Bravina, A. B. Kaidalov and E. E. Zabrodin, J. Phys. G G **35**, 044039 (2008)

[22] V.N. Gribov, Sov. Phys. JETP **29**, 483 (1969); *ibid.* **30**, 709 (1970); *ibid.* **26**, 414 (1968)

[23] A. Schwimmer, Nucl. Phys. B **94**, 445 (1975)

[24] A. Capella, A. Kaidalov, J. Tran Thanh Van, Heavy Ion Phys. **9**, 169 (1999)

[25] N. Armesto, A. Capella, A.B. Kaidalov, J. Lopez-Albacete, C.A. Salgado, Eur. Phys. J. C **29**, 531 (2003)

[26] A. Capella, D. Sousa, Phys. Lett. B **511**, 185 (2001)

[27] A. Capella and E. G. Ferreiro, Eur. Phys. J. C **72**, 1936 (2012)

[28] S. Brodsky, A.H. Mueller, Phys. Lett. B **206**, 685 (1988)

[29] B. Koch, U. Heinz, J. Pitsut, Phys. Lett. **243**, 149 (1990)

[30] A. Capella, Phys. Lett. B **364**, 175 (1995)

[31] A. Capella, A. Kaidalov, A. Kouider Akil, C. Merino, J. Tran Thanh Van, Z. Phys. C **70**, 507 (1996)
A. Capella, C.A. Salgado, D. Sousa, Eur. Phys. J. C **30**, 111 (2003)

[32] B. Abelev *et al.* [ALICE Collaboration], arXiv:1203.3641 [hep-ex].

[33] F. Bossu, Z. C. del Valle, A. de Falco, M. Gagliardi, S. Grigoryan and G. Martinez Garcia, arXiv:1103.2394 [nucl-ex].

[34] B. Abelev [ALICE Collaboration], JHEP **1207**, 191 (2012)

[35] B. Abelev *et al.* [ALICE Collaboration], Phys. Rev. Lett. **109**, 072301 (2012)

[36] J. Wiechula [for the ALICE Collaboration], arXiv:1208.6566 [hep-ex].

[37] E. Scomparin [for the ALICE Collaboration], contribution to Quark Matter 2012;
I. Arsene [for the ALICE Collaboration], contribution to Quark Matter 2012.

[38] R. Arnaldi [for the ALICE Collaboration], contribution to Quark Matter 2012.

[39] A. Rakotozafindrabe, E. G. Ferreiro, F. Fleuret, J. P. Lansberg and N. Matagne, Nucl. Phys. A **855**, 327 (2011)

[40] X. Zhao and R. Rapp, Nucl. Phys. A **859**, 114 (2011)

[41] A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel, J. Phys. G G **38**, 124081 (2011)

[42] E. G. Ferreiro, Contribution to the QCD session of 43rd Rencontres de Moriond 2008 arXiv:0805.2753 [hep-ph].

[43] E. G. Ferreiro, F. Fleuret, J. P. Lansberg and A. Rakotozafindrabe, Contribution to Rencontres de Moriond 2009: QCD and High Energy Interactions, arXiv:0903.4908 [hep-ph].

[44] E. G. Ferreiro, F. Fleuret, J. P. Lansberg and A. Rakotozafindrabe, Phys. Lett. B **680**, 50 (2009)

[45] R. Vogt, Phys. Rev. C **81**, 044903 (2010)

[46] Y. -p. Liu, Z. Qu, N. Xu and P. -f. Zhuang, Phys. Lett. B **678**, 72 (2009)

[47] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1205**, 063 (2012)

[48] C. Mironov [for the CMS Collaboration], contribution to Quark Matter 2012.

[49] D. H. Moon [for the CMS Collaboration], contribution to Quark Matter 2012.